



Ferti-Copper



General Product Use

For use in cases of Increasing Chlorosis, loss of Protein Development, Disease Weakness, Slow Root Activity, Poor Cellular Formation and Lack of Vigour.

Benefits

Copper aids Cell Wall Formation, proper Enzyme Activity and Protein Creation. Promotes several important Chlorophyll functions and is a key Trace Element in Disease Prevention.

First Aid

Follow Acid Protocols.
Harmful if Swallowed.

Wash Eyes. Seek Medical Attention.
If swallowed do not induce vomiting.
Give plenty of water and seek medical attention immediately.



Typical Analysis	% w/v
Copper	6.3
Sulphur	3.2

WITH 0.7% POTASSIUM, 1.9% PHOSPHORUS, FULVIC ACID AND NATURAL CHELATORS

 (08) 9725 6877

PO Box 9129 PICTON WA 6229

www.fertitech.com info@fertitech.com

BATCH NO:

20LT

200LT

1000LT

Directions

AGITATE WELL BEFORE USE

Application Rate

Broadacre

- Apply 0.5-1.5 Litres per Hectare

Orchards, Vines

- Apply 1-3 Litres per Hectare

Horticulture

- Apply 0.5-2 Litres per Hectare

Handling & Safety

Minimise skin contact. Wear Gloves & Eye Protection at all times.

- DO NOT MIX with other products, plant foods etc.

Condition of Sale – Intended Use

This Product is carefully manufactured as fit for its intended purpose. Product Warranty is valid for up to 6 months after Date of Delivery and Product Liability is always subject to the proper use & handling instructions detailed on this label.