



Ferti-IRON™



General Product Use

For use in addressing low soil uptake causing Yield loss. Apply in cases of poor vigour with leaf yellowing and chlorosis. Apply when cold/wet weather affects soil uptake, in higher soil pH (7.5+), in anaerobic soil or where excessive P/Mn exists in soil.

Benefits

Aids Nitrogen Fixation, Essential for Oxygen Carrying, Chlorophyll Production and Enzyme Synthesis. Strongly Enhances Photosynthesis.

First Aid

Harmful if Swallowed.

Follow Acid Hazard Protocols.

Wash Eyes. Seek Medical Attention.

If swallowed do not induce vomiting. Give plenty of water and seek medical attention immediately.



Typical Analysis % w/v

Iron (EDTA) 6.2

EDTA-CHELATE BUFFERED WITH HIGH LEVELS OF FERTI-FULVIC ACID FOR A MORE SUITABLE AND EFFECTIVE CELLULAR TRANSLOCATION

Directions

AGITATE WELL BEFORE USE

Application Rate

Broadacre

- Apply 2-4 Litres per Hectare

Orchards, Vines

- Apply 2-5 Litres per Hectare

Horticulture

- Apply 3-6 Litres per Hectare

Handling & Safety

Minimise skin contact. Wear Gloves & Eye Protection at all times.

- DO NOT MIX with other products, plant foods, pesticides etc.

Condition of Sale – Intended Use

This product is carefully manufactured as fit for its intended purpose.

Product Warranty and Liability is always subject to the proper use and handling instructions detailed on this label.



(08) 9725 6877

PO Box 9129 PICTON, WA 6229

www.fertitech.com info@fertitech.com

BATCH NO:

20LT

200LT

1000LT